

Spatial Relationships of Plant Bugs in Large Scale Cotton Operations: Do Edge and Ecotone Matter?

Isaac L. Esquivel^{1,2}, Michael J. Brewer^{1,2}, Robert N. Coulson¹

¹Texas A&M University Department of Entomology, College Station ²Texas A&M AgriLife Research and Extension Center, Corpus Christi

ABSTRACT

The Cotton Flea Hopper, and the Verde Plant bug (Hemiptera: Miridae) have become more problematic in recent years in the cotton growing region of South Texas. They feed on reproductive tissue: squares (cotton flea hopper) and young bolls (verde plant bug). Feeding during these stages can cause severe damage to the final cotton yield as well as introduce cotton boll rot. A pest management issue is that outbreaks vary from year to year and their densities vary from field to field. Edges and ecotones are often used in landscape ecology to aid in interpreting the functional heterogeneity of an organism. In the case of these cotton insects, how the surrounding landscape effects their population occurrence and movement from squaring through flowering. The goal of this study was to investigate the spatial relationships with these pest populations to get a better understanding on why some of these pests are present in some areas and not present in others. Our study found both a edge effect for both verde plant bugs and cotton flea hoppers (more bugs were found nearer to the field edge) and ecotone modifies the strength of the edge effect. Edge effect was more predominant in the Cotton/Natural habitat ecotone than the others. This information can prove to be useful in pest management as it may help managers focus on fields or areas of fields that may be of higher risk to colonization by these pests.

OBJECTIVES

- Investigate an edge effect on the population of cotton flea hopper and verde plant bug.
- Investigate the effect of ecotone on the populations of cotton flea hopper and verde plant bug and identify any interaction between ecotone and edge on populations of these pests.

INTRODUCTION

The discipline of landscape ecology focuses on how landscape patterns, biological entities, ecological processes interact. It serves to explore these interactions within a spatial context.¹

The cotton flea hopper (Figure 1a) feeds on cotton squares. The verde plant bug (Figure 1b) on the other hand tends to feed on young bolls. These pests have one aspect in common, they both need an overwintering site after crops have been harvested and destroyed. The flea hopper tends to overwinter on woolly croton and selected other plants. The verde plant bug uses *Saueda* spp., a succulent type plant for overwintering. Both of these host plants occur throughout the mixture of grass and shrub land in natural habitat. Many agricultural fields are juxtaposed on this natural habitat matrix.




Figure 1c: Visual interpretation of edge vs ecotone. In this case 'edge' is the hard line labeled border but as you can see there is a transition between the two land cover types in the image. KEL image adapted from Forman 1995

Edges are defined as the distinct border between two different land cover types and can be hard or soft, curvilinear or straight, and natural or anthropogenic. In our case, the cover types are crops (cotton and sorghum) and are anthropogenic edges and natural habitat that borders the fields. An edge effect refers to the high population density and diversity of a species or species in the outer portion or edge of a patch of habitat, in this case fields. Ecotones are defined as the area occurring at the interface of two or more distinct landscape elements (Figure 1c). In this study these would be the transition between different crops (cotton-cotton or cotton-sorghum) and natural habitat. An investigation of these effects can give us a better understanding at the spatial dynamics of these pests and perhaps provide useful information for pest management strategies.



Figure 1a: Cotton Flea Hopper courtesy of bugguide.

Figure 1b: Verde Plant Bug courtesy of Bugguide.

STUDY AREA

Our study area is a large commercial cotton and grain farming system in the coastal cotton growing region of South Texas (Fig.2). It is about a 30,000 acres system of cotton, sorghum, and unmanaged habitat. The unmanaged habitat is composed of mixed grass/shrub land and a bay/estuary zone that penetrates into the cropping system. There are many overwintering sites of our primary study insect pests, as well as beneficial organisms




Figure 2: Graphic of cotton region represented of the study area. Created by I.E. Esquivel

METHODS

Since our research area provides a unique environment of large scale agricultural fields juxtaposed along natural habitat areas, it provides a set up to investigate edge and ecotone effects.

Objective 1: Edge effects

- Fields were selected based on the diversity of the surrounding fields and natural habitat area surrounding them.
- A series of parallel transects 100 meters apart were constructed using ESRI's ArcMap gis software then imported into a Trimble Juno 3b GPS unit running ArcPad. This allowed efficient collection of geo-reference data.
- Using the Juno 3b GPS system, insect samples were taken at the specified localities at the edge 0m, 10m, 25m, 100m, 200m and 300m into the field (Figure 3).
- For Cotton Flea Hopper; 40 plants were sampled using the beat bucket method and 20 plants were inspected for square damage.
- For Verde Plant Bug; 40 plants were sampled using the beat bucket method and 20 quarter sized green bolls were collected and inspected in the lab for internal damage.

Objective 2: Ecotone effects

- Ecotones of interest are cotton adjacent to cotton (CC), cotton adjacent to sorghum (CS) and cotton adjacent to natural habitat (CN).
- Two fields gave us all three transitions allowing us to proceed with transect sampling in each ecotone.
- Ecotones were used as a covariate (CC,CN or CS) in analyzing the transects to investigate the ecotone effect and any interaction ecotone has on pop density.



Transect Setup




Figure 3: Example of one transect along one edge of a cotton field chosen. This transect depicts the cotton-natural habitat ecotone where the cotton field abuts the natural habitat.

RESULTS: Flea Hopper

NOTE ON MAPS: There are two fields of interest (Field 1&2) will be labeled in the captions. Circles represent pest densities per plant and are represented by graduated color symbols increasing from 0. Cutoffs represent the economic thresholds (orange and red dots).




Figure 4: Transects of field 1 showing the cotton-cotton, cotton-sorghum, and cotton-natural habitat ecotones.

Results Continued...




Figure 4a: Zoomed in view from fig 4 of field 1, transect showing a significant effect of distance from edge on flea hopper density.




Figure 4b: Linear regression of flea density vs distance. R²: .5

Edge effect?

When all cotton flea hopper densities were plotted against distance from edge, there was a significant negative relationship ($p < .0001$) however there was a weak R² value of 0.18. When densities were plotted against distance by the Cotton-Natural habitat ecotone, it strengthened the negative relationship with a much improved R² of 0.5 (Fig. 4a.b.). This is similar to our analysis of distance with ecotone as a covariate (Fig. 5).

Ecotone effect?

Flea densities were plotted against distance with ecotone as a covariate to test for an ecotone effect. There was a significant distance from edge effect ($p < .001$) and a significant ecotone effect ($p < .001$). There also appears to be a significant interaction between ecotone and distance ($p < .001$), which indicates flea hopper densities on the edges are influenced by the ecotone. Overall the covariate regression model was significant with an R² value of 0.4 (Fig.5).




Figure 5: Transects of field 2 showing the cotton-cotton, cotton-sorghum, and cotton-natural habitat ecotone (left to right).

RESULTS: Verde Plant Bug




Figure 6: Transect from field 1 showing a significant edge effect for verde plant bugs at the cotton-natural habitat ecotone.

Results Continued...




Figure 7: Visualization of Verde plant bugs densities in field 2 with all three ecotones present: Cotton-Cotton, Cotton-sorghum and Cotton-Natural Habitat. The right transect is the one of interest, appears to have more insects than others alongside the natural habitat.




Figure 8a: Linear regression of Verde density vs distance. R²: .5

Edge Effect?

Verde plant bug densities were plotted against distance from edge, there was a significant negative linear relationship ($p < .001$) however there was a weak R² value of 0.11. When the densities were plotted individually by 'ecotone' there were also significant linear relationships. For example fig 8a shows the relationship between Verde density and distance for values in the cotton-natural habitat ecotone with an R² of 0.5. Similar results were found when analyzing an ecotone effect (fig. 8b).

Ecotone Effect?

Verde plant bug densities were plotted against distance from the edge with ecotone as a covariate to test for an ecotone effect. In our covariate analysis there was a significant edge effect with a ($p < .0001$) and a significant ecotone effect with a ($p < .0001$). There was also a significant interaction between ecotone and Distance ($p .0291$). This would suggest that much like flea hoppers, there appears to be more verde plant bugs on average at the edge of the cotton-natural habitat ecotone.

DISCUSSION

Objective 1:

Analysis of an edge effect for flea hopper indicates that overall there is a significant negative relationship with distance. However, when broken down by ecotone we also see a significant relationship with flea density with a much improved R² of 0.5. This would suggest that there is a weak edge effect overall but certain areas of the field have a strong edge effect while others may have a weaker or positive edge effect like that for flea hoppers. The same phenomena happens with respect to Verde plant bugs. When analyzed as a whole, there was a significant linear relationship but a weak R². However when plotted separately by ecotone there was also a significant negative relationship with a much improved R² value of 0.5 much like flea hopper.

Objective 2:

When analyzing the same data with ecotone as a covariate we see further evidence that ecotone has an effect on both the cotton flea hopper and verde plant bug densities. Our covariate analysis shows significant distance and ecotone effects. However, there is also a significant interaction between ecotone and distance. This supports that distance from the edge has a significant effect on flea hopper and verde plant bug densities but only when you take into account the ecotone. For example, the edge effect on the interface of cotton and natural habitat is statistically different from the edge effect at the interface of cotton and sorghum. These effects are driven by the cotton-natural habitat ecotone (fig 7) in which on average, there are more verdes and fleas at edge of the field in that ecotone compared to the cotton-cotton and cotton-sorghum ecotone.

Our findings can be useful for managers of large scale cotton systems as it appears pest densities are higher on average on sides of the field that abut natural habitat and tend to congregate at the edge of the fields between 0m - 100m. This may help save time and focus areas for scouting.

ACKNOWLEDGMENTS

Funding by the Everette C Salyer Fellowship. I would like to thank members of the Knowledge Engineering lab for their helpful comments and suggestions when planning this work. I would like to thank members of the field crop entomology lab at the AgriLife Research lab in Corpus Christi including: Leo DeLeon, Micheal Aldridge, Travis Aherns, and Luke Pruter for help with data collection in the field. Also, Mike Brewer and Darwin Anderson for help in planning.

REFERENCES

¹Coulson, R. and M. Tolkeren. 2010. Basic landscape ecology. KEL, College Station, TX.
²Paraglider, M. N., et al. 2008. J. Econ. Entomol. 99: 268-277. Paper covers: sampling methods and dispersion for western flower thrips and cotton fleahoppers in cotton.
³Brewer, M. J., et al. 2012. J. Econ. Entomol. 105: 898-905. Paper covers: sampling strategies for cotton fleahopper and verde plant bug occurring on cotton.